

Confinement/Deconfinement Transition of Large N Gauge Theories in Perturbation Theory with N_f Fundamentals: N_f/N Finite

Howard J. Schnitzer¹
Theoretical Physics Group
Martin Fisher School of Physics, Brandeis University
Waltham, MA 02454

Abstract

In hep-th/0402219 we considered large N zero-coupling d -dimensional $U(N)$ gauge theories, with N_f matter fields in the fundamental representation on a compact spatial manifold $S^{d-1} \times \text{time}$, with N_f/N finite. This class of theories undergo a 3rd order deconfinement transition. As a consequence it was proposed that the dual string theory has a 3rd order phase-transition to a black hole at high temperature.

In this paper we argue that the same conclusions are valid for such theories at any finite order in perturbation theory in the 't Hooft coupling λ and N_f/N . It is conjectured that this continues to hold at strong coupling and finite value of $N_f/N < 1/\lambda$, which suggests that the supergravity approximation to the dual string theory has a 3rd order thermal phase transition to a large black hole.

1 Introduction

The consideration of large N gauge theories, their confinement/deconfinement phase-transitions and the possible relation to the transition from a thermal ensemble to a large black hole at high-temperatures in the dual string theory is a topic which is being actively discussed [1]. An important insight [2]–[4] was made by considering the gauge theories on a compact space, as there is an additional parameter $R\Lambda$ which can be tuned to weak-coupling allowing for a perturbation expansion, where R is the size of the compact space, and Λ is the dynamical scale of the gauge theory. As the gauge theory is on a compact manifold, the Gauss law constraint induces interactions between gluons and the matter multiplets even at zero coupling.

Sundborg [2] and Aharony, *et al.* [3, 4], have provided a general framework to discuss these issues in the context of a unitary matrix model. This strategy has been developed and extended in a variety of ways [5]–[8]. The particular system that concerns us in this paper is large N , $U(N)$ gauge theory with N_f matter multiplets in the fundamental representation with N_f/N finite.

We previously have shown [5] that large N , zero-coupling, $U(N)$ gauge theory with N_f matter fields in the fundamental representation on a compact spatial manifold $S^{d-1} \times S^1$, with (N_f/N) finite, undergoes a 3rd order confinement/deconfinement phase transition at a temperature T_c . The

¹email: schnitzr@brandeis.edu

Supported in part by the DOE under grant DE-FG02-92ER40706

low-temperature phase (phase A_0) has free energy of $\mathcal{O}(N_f^2)$, interpreted as a gas of color singlet mesons and glueballs, while the high-temperature phase (phase A_1) has a free energy characterized by $N^2 f(N_f/N, T)$, interpreted as that of a gas of gluons and of fundamental and anti-fundamental matter states.

An intriguing question is the identification of the string dual to the gauge theory and the associated deconfinement transition. Witten [1] and others [2]–[8] have associated the confinement/deconfinement transition with the Hawking–Page [9] transition of the dual theory. When ($N_f = 0$) the low-temperature phase has free energy of $\mathcal{O}(1)$, while the high temperature phase has free energy of $\mathcal{O}(N^2)$, signaling the deconfinement of color singlet glueballs into gluons. In the dual string theory this is associated to a transition from a thermal gas of supergravity modes to a large black hole [1]. At zero-coupling of the gauge theory on $S^{d-1} \times S^1$, this transition is 1st order at the Hagedorn temperature T_H [2, 3]. For pure Yang–Mills theory on $S^3 \times S^1$, computed to 3-loops in perturbation this remains a first-order transition [3, 4]. At weak-coupling in the gauge theory, one expects the thermal string/black hole transition to occur for black holes with size of order of the string scale, while a comparison with the supergravity approximation requires strong-coupling in the gauge theory.

When (N_f/N) is finite, at zero-coupling the transition is 3rd order on the gauge side [5]. However, no example is yet known of a string theory with a 3rd order transition to a black hole. The challenge is to understand if such a transition exists in the corresponding string dual. If the zero-coupling results of a 3rd order transition survives extrapolation to strong-coupling, then one would have the (relatively) easier task of finding an appropriate dual in the supergravity approximation. This task itself is not without difficulties. For example, with $d = 4$ one might be tempted to consider a IIB model with N $D3$ branes and N_f $D7$ branes, with (N_f/N) finite in the large N limit. However, this simple picture is not possible as the number of allowed $D7$ branes is limited. One approach to this issue is to consider $d=3$, with N $D2$ branes and N_f $D6$ branes at finite temperature, including back-reaction [10]. Unfortunately the construction considered in ref. [10] only gave the high-temperature limit of the system, so there was no opportunity to observe any possible Hawking–Page transition. Another approach does allow for a back reaction in a $d=4$ string theory with $N_f \neq 0$ [11], but again the putative phase-transition has not been discussed. Finally, if one has N_f probe $D7$ branes embedded in $AdS_5 \times S^5$ [12, 14], the Hawking–Page transition remains first-order even at strong-coupling but in the probe approximation [14, 6]. Perhaps this is not surprising as $N_f/N \rightarrow 0$ in the large N limit in the probe approximation, so that this should be compatible with $N_f = 0$. Therefore the possibility of a 3rd order Hawking–Page transition with N_f/N finite is unresolved.

In this paper we continue the discussion began in ref. [5] where N_f/N is finite, but now with non-vanishing coupling constants. In Sec. 2 we consider the planar contribution at large N to S_{eff} (the effective action) in perturbation theory, as a double expansion in the 't Hooft coupling λ and N_f/N . This is presented in terms of a matrix model for U , given in terms of the zero-mode of the gauge field

$$\alpha = \frac{1}{\omega_{d-1}} \int_{S^{d-1}} A_0 , \quad (1.1)$$

$$U = e^{i\beta\alpha} = e^{i\alpha/T} \quad (1.2)$$

and

$$u_n = \frac{1}{N} \text{Tr}(U^n) . \quad (1.3)$$

We argue in Sec. 2 that at low-orders in the double perturbation theory, there is no solution to the saddle-point equations with $u_n = 0$ ($n \geq 1$). Since the eigenvalue distributions of U can be written as

$$\rho(\theta) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} u_n e^{in\theta} \quad (1.4)$$

where $u_n = \bar{u}_n$ and $u_0 = 1$, thus there is *no* phase with $\rho(\theta) = \text{constant}$.

In Sec. 3 we discuss in detail the phase structure as a function of temperature that follows from the S_{eff} of Sec. 2 for the special case where $(N_f/N) = \mathcal{O}(\lambda^2)$. In this example the saddle-point equations require solution of cubic equations, and already exhibit many of the main issues of this paper. There is a 3rd order transition from phase A_0 to A_1 [5, 13] at a temperature T_c below the putative Hagedorn temperature, a feature which appears to be a generic property of gauge theories with (N_f/N) finite.

In Sec. 4 we adapt the strategy of ref. [8] to the issues of this paper. In eq'n. (4.1) we present the gauge invariant matrix model consistent with the double perturbation expansion in λ and (N_f/N) of large N non-Abelian $U(N)$ gauge theory. First we recover the zero-coupling results of ref. [5]. The saddle-point equations which follow from (4.1) are shown in general to exclude the phase described by $\rho(\theta) = \text{constant}$, to any finite order in the double perturbation series in λ and (N_f/N) . Thus in this case, the low-temperature phase is always in phase A_0 , *i.e.*, that described by $\rho(\theta) \neq \text{constant}$, covering $0 \leq \theta < 2\pi$. We argue, again in perturbation theory, that the transition to the single-gapped phase A_1 is 3rd order.

Sec. 5 summarizes our results, and offers plausible conclusions regarding the string dual evaluated in the supergravity approximation. It is reasonable to expect that the supergravity limit of string theories dual to the class of gauge theories considered in this paper exhibit a 3rd order transition from a thermal gas of supergravity modes to a large black hole. No such example is yet known.

Issues not discussed in this paper are:

1. small black-holes [8, 16],
2. smoothing out the phase-transitions at finite N [7, 8], and
3. the logical possibility that at a temperature above T_c , there exist phases A_n such that $\rho(\theta)$ is n -gapped [15].

Since even when (N_f/N) is $\mathcal{O}(\lambda^2)$, *c.f.* Sec. 3, the solution of the saddle-point equations require analysis of a cubic, the general analysis of the saddle-point equations can involve complicated, model-dependent algebraic equations. The principle conclusions of this paper do not depend on such considerations.

2 The Effective Action

A. Perturbation theory

Consider free $U(N)$ gauge theories on $S^{d-1} \times \text{time}$, with N_f matter multiplets in the fundamental representation of the gauge group with N_f/N finite as $N \rightarrow \infty$. The partition function on this compact space (therefore subject to Gauss' law constraints) is a matrix model

$$\begin{aligned} \mathbb{Z}(x) = & \int [dU] \exp \left\{ \sum_{n=1}^{\infty} \frac{1}{n} \left[\mathcal{Z}_B(x^n) + (-1)^{n+1} \mathcal{Z}_F(x^n) \right] \cdot \left[\text{tr}(U^n) + \text{tr}(U^{+n}) \right] \right. \\ & \left. + \sum_{n=1}^{\infty} \frac{1}{n} \left[z_B(x^n) + (-1)^{n+1} z_F(x^n) \right] \text{tr}(U^n) \text{tr}(U^{+n}) \right\} \end{aligned} \quad (2.1)$$

where $z_B(x)$ and $z_F(x)$ are the single-particle partition functions for the bosons and fermions in the adjoint representation, while $\mathcal{Z}_B(x)$ and $\mathcal{Z}_F(x)$ are the single-particle partition functions for the bosons and fermions in the fundamental representation, where

$$x = e^{-\beta} = e^{-1/T} \quad (2.2)$$

for S^{d-1} with radius $R=1$. [R may be restored from dimensional considerations as needed.] Recall that

$$u_n = \frac{1}{N} \text{Tr}(U^n) \quad (2.3)$$

Now consider the same theory with non-vanishing gauge couplings g^2 , with $\lambda = g^2 N$ the 't Hooft coupling. It is useful, in this section without a specific theory in mind to keep any scalar self-couplings and Yukawa couplings at zero. The partition function can be written as

$$\mathbb{Z}(x) = \int du_n d\bar{u}_n \exp \left[-\tilde{S}_{eff}(u_n) \right]. \quad (2.4)$$

The planar contribution at large N to S_{eff} in perturbation theory, in a double expansion in λ and N_f/N , consistent with gauge invariance, is

$$\begin{aligned} N^{-2} \tilde{S}_{eff}(u_n) = & S_{eff}(u_n) \\ = & \left\{ \sum_n G_n(x) |u_n|^2 \right. \\ & + \lambda \sum_{m,n} G_{m,n}(x) (u_m u_n u_{-n-m} + c.c.) \\ & + \lambda^2 \sum_{m,n,p} G_{m,n,p}(x) (u_m u_n u_p u_{-n-m-p} + c.c.) \\ & + \dots \left. \right\} \\ & + \left(\frac{N_f}{N} \right) \left\{ \sum_n F_n(x) [u_n + c.c.] + \lambda \sum_{m,n} F_{m,n} [u_n u_m u_{-n-m} + c.c.] \right. \\ & + \lambda^2 \sum_{n,m,p} F_{n,m,p}(x) [u_n u_m u_p u_{-n-m-p} + c.c.] \\ & + \lambda^3 \sum_{n,m,q,p} F_{n,m,q,p}(x) [u_n u_m u_p u_q u_{-n-m-p-q} + c.c.] \\ & \left. + \dots \right\} + \mathcal{O} \left(\left(\frac{N_f}{N} \right)^2 \right). \end{aligned} \quad (2.5)$$

Consider the Feynman diagrams of the theory, where each insertion of a fundamental loop into an adjoint line of a diagram gives a factor of $\lambda(N_f/N)$, while a radiative correction by an adjoint line to a fundamental propagator gives a factor of λ . [Since we are keeping scalar self-couplings and Yukawa couplings at zero, these are the only corrections to be considered in this section.] In (2.5) the leading powers of λ and of (N_f/N) in the expansion of $\tilde{S}_{eff}$ are exhibited explicitly. The various coefficients in (2.5) are a double power series in λ and (N_f/N) , beginning at $\lambda = 0$ and $(N_f/N) = 0$.

Near the saddle point of (2.5) one may integrate out the $u_n (n \geq 2)$, since at the critical point x_c in phase A_0 , $|u_n(x_c)| \ll |u_1(x_c)|$ for $n \geq 2$. Therefore the effective action for u_1 near the saddle-point for u_1 is

$$\begin{aligned}
N^{-2} \tilde{S}_{eff}(u_1) &= \left[A|u_1|^2 + B|u_1|^4 + \dots \right] \\
&+ \left(\frac{N_f}{N} \right) \left\{ (u_1 + \bar{u}_1) \left[-C + D|u_1|^2 + E|u_1|^4 + \dots \right] \right. \\
&+ \left(u_1^2 + \bar{u}_1^2 \right) \left[F + G|u_1|^2 + H|u_1|^4 + \dots \right] \\
&+ \left(u_1^3 + \bar{u}_1^3 \right) \left[K + L|u_1|^2 + M|u_1|^4 + \dots \right] \Big\} \\
&+ \dots
\end{aligned} \tag{2.6}$$

The leading order contributions to the coefficients in (2.6) are

$$\begin{aligned}
A &= \mathcal{O}(1) ; B = \mathcal{O}(\lambda^2) ; \\
C &= \mathcal{O}(1) \text{ and positive at leading order} ; \\
D &= \mathcal{O}(\lambda^2) ; E = \mathcal{O}(\lambda^4) ; \\
F &= \mathcal{O}(\lambda) ; G = \mathcal{O}(\lambda^3) ; H = \mathcal{O}(\lambda^5) ; \\
K &= \mathcal{O}(\lambda^2) ; \text{etc.}
\end{aligned} \tag{2.7}$$

Since each insertion of a fundamental loop in an adjoint line is accompanied by a factor of $\lambda(N_f/N)$, higher-orders in (N_f/N) result in higher powers of λ compared to those displayed in (2.6), while the $\mathcal{O}(N_f/N)$ term is that of a single fundamental loop with adjoint radiative corrections. Consider the schematic structure of the saddle-point condition which follows from (2.6)

$$\begin{aligned}
&N^{-2} \frac{\partial \tilde{S}_{eff}}{\partial u_1} \\
&= - \left(\frac{N_f}{N} \right) C + u_1 \left\{ 2B \bar{u}_1^2 + 2 \left(\frac{N_f}{N} \right) F + \dots \right\} \\
&\quad + \bar{u}_1 \{ A + \dots \} + \mathcal{O} \left(\lambda \left(\frac{N_f}{N} \right)^2 \right) = 0
\end{aligned} \tag{2.8}$$

where the curly brackets are a power-series in u_1 and $\bar{u}_1$, with coefficients a series in λ . [We have only exhibited the leading terms explicitly in the curly brackets.] Recall that C is a power series in λ , with its leading term $\mathcal{O}(1)$ real and positive, while A has leading-term $\mathcal{O}(1)$, F of $\mathcal{O}(\lambda)$, and B of $\mathcal{O}(\lambda^2)$. Thus, $u_1 = 0$ is *not* a solution to (2.8) to the order we are working. Further, since corrections to the constant term in (2.8) are a power series in λ and $(\lambda N_f/N)$, the general structure of the saddle-point equation in perturbation theory, emphasizing the constant terms, is

$$\begin{aligned}
& N^{-2} \frac{\partial \tilde{S}_{eff}}{\partial u_1} \\
&= - \left(\frac{N_f}{N} \right) C + \sum_{n=1, m=0}^{\infty} \lambda^{n+m} \left(\frac{N_f}{N} \right)^{n+1} C_{n,m} \\
&\quad + u_1 \left\{ 2B \bar{u}_1^2 + \dots \right\} \\
&\quad + \bar{u}_1 \left\{ A + \dots \right\} = 0 .
\end{aligned} \tag{2.9}$$

The curly brackets represent a perturbative expansion in u_1 with the leading term exhibited explicitly, where the C_n are independent of u_1 and $\bar{u}_1$, but a power-series in λ . Since $C = \mathcal{O}(1)$, and positive at leading order, *cf.* (2.7), $u_1 = 0$ is *not* a solution to (2.9) in any finite order of the double perturbation theory, as the powers of λ and (N_f/N) do not match. A solution $u_1 = 0$ to (2.9) would require

$$\sum_{n=1, m=0}^{\infty} \lambda^{n+m} \left(\frac{N_f}{N} \right)^n C_{n,m} \stackrel{?}{=} C \tag{2.10}$$

which is not possible in any finite order of the double expansion, since C at leading order is independent of λ and (N_f/N) . If (2.10) has a solution, it must be a non-perturbative or all-orders result. As a consequence, $u_1 = 0$ does *not* solve the saddle-point equations in perturbation theory if (N_f/N) finite is considered as an expansion parameter of the theory.

B. Phases

The eigenvalue distribution of the matrix U can be written as

$$\rho(\theta) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} u_n e^{in\theta} \tag{2.11}$$

where $u_{-n} = \bar{u}_n$ and $u_0 = 1$. The u_n , with $n \geq 2$ can be expressed in terms of u_1 near the saddle-point by means of the saddle-point equations that follow from (2.5), which is how (2.6) is obtained. Schematically, (not being explicit about coefficients and not distinguishing between u_n and u_{-n}), one has

$$\begin{aligned}
u_2 &\sim \left(\frac{N_f}{N} \right) + \lambda \left(\frac{N_f}{N} \right) u_1 + \lambda u_1^2 + \dots \\
u_3 &\sim \left(\frac{N_f}{N} \right) + \lambda u_1 u_2 + \lambda \left(\frac{N_f}{N} \right) u_1 + \dots \\
&\sim \left(\frac{N_f}{N} \right) + \lambda \left(\frac{N_f}{N} \right) u_1 + \lambda^2 \left(\frac{N_f}{N} \right) u_1^2 \\
&\quad + \lambda^2 u_1^3 + \dots ,
\end{aligned} \tag{2.12}$$

which displays the leading orders in λ and (N_f/N) in (2.12), and similarly with its generalization to u_n . Eqn. (2.12) reiterates that $u_n \neq 0$ are the only solutions to the saddle-point equations within the context of perturbation theory.

Thus, for (N_f/N) finite in the large N limit, (2.11)–(2.12) and the saddle-point equations require, in the double perturbation expansion that

$$\rho(\theta) \neq \text{constant} . \tag{2.13}$$

Therefore, the allowed phases (in the notation of ref. [5]) are

- (i) $(A_0) : \rho(\theta)$ completely covers the circle, but is not constant. (2.14)
- (ii) $(A_1) : \rho(\theta) = 0$

for a single finite gap in the distribution. The phase (A_0) is that described by the saddle-point, while (A_1) can be described by the methods of refs. [2, 3, 5, 8, 13]. The criterion that $\rho(\theta) \geq 0$ implies from (2.12)–(2.13) that

$$|u_1| \leq \frac{1}{2} + \mathcal{O}\left(\frac{N_f}{N}\right) + \mathcal{O}(\lambda) + \mathcal{O}\left(\lambda \frac{N_f}{N}\right) + \dots \quad (2.15)$$

in the double perturbation theory. The (A_0, A_1) phase boundary occurs when $\rho(\theta) = 0$ at a single point, which then implies that (2.15) is an equality. The transition to phase A_1 is third-order, as can be deduced from an analysis analogous to that of ref. [5]. [It is a logical possibility [15] that there could be additional phases A_n , with n finite gaps ($n \geq 2$), whose phase boundary occurs at temperatures above that of the (A_0, A_1) transition, but this is outside the scope of our discussion. Further, this does not occur in theories with $\lambda = 0$, $(N_f/N) \neq 0$ [5].]

3 Phase Structure of an Example

In this section we discuss in detail the phase structure for the case when $(N_f/N) = \mathcal{O}(\lambda^2)$. The purpose of the example is to illustrate a detailed discussion of the saddle-point equations, and evolution of the phase-structure with temperature. Although there are only two phases in this case, the details of the evolution with temperature depend on the relative magnitudes of parameters. When $(N_f/N) = \mathcal{O}(\lambda^2)$, insertions of fundamental loops into the diagrams of pure Yang–Mills theory can be neglected to the order in λ^2 we are considering. Further, this example is interesting, as one can additively combine the contribution of fundamental matter to $\tilde{S}_{eff}$ with the results of ref. [4], where pure Yang–Mills theory in $d=4$ was analyzed to three-loop order.

A. General considerations

Consider (2.6)–(2.7) correct to $\mathcal{O}(\lambda^2)$, with $(N_f/N) = \mathcal{O}(\lambda^2)$. Only the first three terms of (2.6) have coefficients of that order so that to leading order

$$\begin{aligned} N^{-2} \tilde{S}_{eff}(u_1) &= m_1^2 |u_1|^2 + b |u_1|^4 \\ &\quad - d(u_1 + \bar{u}_1) \end{aligned} \quad (3.1)$$

where

$$m_1^2 = K(T_H - T) ,$$

with

$$K = \mathcal{O}(1) , \quad (3.2)$$

$b = \mathcal{O}(\lambda^2)$, $d > 0$ and $\mathcal{O}(N_f/N)$, $(N_f/N) = \mathcal{O}(\lambda^2)$, and where T_H is defined as the temperature where $m_1^2 = 0$. Separate $u_1 = \tilde{u}_1 + i\tilde{u}_1$ into real and imaginary parts, so that the saddle-point equations that follow from (3.1) are

$$\begin{aligned} m_1^2 \tilde{u}_1 + 2b \tilde{u}_1 |u_1|^2 - d &= 0 \\ m_1^2 \hat{u}_1 + 2b \hat{u}_1 |u_1|^2 &= 0 \end{aligned} \quad (3.3)$$

which requires $\hat{u}_1 = 0$; consistent with $d = (N_f/N)C$ real. To the order we are considering, C is computed at $\lambda = 0$, so that it has the same value as in ref. [5]. The restriction (2.15) now becomes $|u_1| \leq 1/2 + \mathcal{O}(\lambda)$ in phase A_0 .

In order to study phase A_0 , we need the saddle-point solution to

$$N^{-2} \tilde{S}_{eff}(u_1) = m_1^2 |u_1|^2 + b |u_1|^4 - 2 d u_1 \quad (3.4)$$

which requires

$$2b u_1^3 + m_1^2 u_1 - d = 0. \quad (3.5)$$

Solutions to (3.5) are characterized by the discriminant

$$D = -\frac{1}{2} \left(\frac{m_1^2}{b} \right)^3 - 27 \left(\frac{d}{2b} \right)^2. \quad (3.6)$$

When $D > 0$ there are 3 real roots from (3.5), while if $D < 0$ there is 1 real root, and 2 complex roots which are not relevant, as u_1 is real. If

$$\left| \frac{d}{2b} \right|^{2/3} \gg \left| \frac{m_1^2}{3b} \right| \quad (3.7)$$

then $D < 0$.

If

$$\left| \frac{d}{2b} \right|^{2/3} \ll \left| \frac{m_1^2}{3b} \right|, \quad (3.8)$$

the sign of D depends on the sign of (m_1^2/b) , in which case the solution for u_1 is approximately that of ref. [3, 4] *except* instead of $u_1 = 0$, now

$$u_1 \simeq (d/m_1^2) > 0 \quad \text{for } m_1^2 > 0.$$

Since $(d/2b) \sim \mathcal{O}(1)$ in this section, (3.8) always occurs for T sufficiently below T_H . For example, $m_1^2 \sim \mathcal{O}(\lambda)$ is sufficient for this to occur.

In the next subsection we concentrate on a toy model, with $b < 0$, recalling that $b < 0$ for pure Yang–Mills theory in $d=4$ [3]. The analysis for $b > 0$ is qualitatively similar, which we do not present here.

B. Toy model

In the general case, $u_n \neq 0$ for $n \geq 1$, which leads to the constraint (2.15). In this section $(N_f/N) \sim \mathcal{O}(\lambda^2)$, so that

$$|u_1| \leq 1/2 + \mathcal{O}(\lambda)$$

in the saddle-point phase A_0 . One can simplify the analysis without sacrificing any of the qualitative features of the physics, so that for the remainder of this section we consider (3.4), but with $u_n = 0$ for $n \geq 2$. Then $\rho(\theta) = 1/2\pi (1 + 2u_1 \cos \theta)$, which implies that

$$|u_1| \leq 1/2 \quad (3.9)$$

with the (A_0, A_1) phase boundary occurring when $|u_1|$ increases to $|u_1| = 1/2$ in the A_0 phase, with the transition to A_1 of 3rd order.

C. $b < 0$

We track the evolution of the system for increasing T , beginning with $T < T_H$.

(i) $m_1^2 > 0$; $D > 0$

If $D > 0$, this requires

$$|b| < \frac{2}{27} \frac{(m_1^2)^3}{d^2} \quad (3.10)$$

and an approximate solution to (3.5) is

$$\begin{aligned} u_1 &\simeq d/m_1^2 \\ &< \left[\frac{2}{27} \frac{d}{|b|} \right]^{1/3}. \end{aligned} \quad (3.11)$$

[The two other real solutions to (3.5) are that of maxima of $\tilde{S}_{eff}(u_1)$, and thus not relevant.] For sufficiently large m_1^2 [$m_1^2 \sim \mathcal{O}(\lambda)$ is sufficient], $|u_1| < 1/2$ so that the system is in phase A_0 , and

$$N^{-2} \tilde{S}_{eff}(u_1) \simeq -d^2/m_1^2. \quad (3.12)$$

The free-energy is

$$F/T = \tilde{S}_{eff}(u_1) \quad (3.13)$$

so that

$$\begin{aligned} F/T &= -N^2(d^2/m_1^2) \\ &\simeq \mathcal{O}(N_f^2)/(T_H - T), \end{aligned} \quad (3.14)$$

since $d \sim \mathcal{O}(N_f/N)$. Thus (3.14) has the same form as (1.1) of ref. [5], and the interpretation of phase A_0 as that of a gas of gluons and color singlet mesons. Therefore for sufficiently low temperatures one is in phase A_0 .

The saddle-point value of u_1 increases as T increases. Consider

(ii) $D > 0$, near $D = 0$; $m_1^2 > 0$

Using (3.6) we have

$$m_1^2 = \left[\frac{27}{2} d^2 |b| \right]^{1/3} + \delta m^2, \quad (3.15)$$

and

$$\begin{aligned} u_1 &= \left[\frac{d}{4|b|} \right]^{1/3} \left\{ 1 - \mathcal{O} \left(\frac{\delta m^2}{m_1^2} \right) \right\} \\ &= \frac{d}{2m_1^2} \left\{ 1 - \mathcal{O} \left(\frac{\delta m^2}{m_1^2} \right) \right\}. \end{aligned} \quad (3.16)$$

We verify that $\frac{\partial^2 \tilde{S}_{eff}}{\partial^2 u_1}(u_1) > 0$, so that this minimum is stable.

If $u_1 < 1/2$, the system is still in phase A_0 at this temperature, while if $u_1 > 1/2$ the system has already made a transition to phase A_1 at a lower-temperature.

(iii) $D = 0$; $m_1^2 > 0$

$$\begin{aligned} u_1 &= \left[\frac{d}{4|b|} \right]^{1/3} \\ &= \frac{d}{2(m_1^2)} . \end{aligned} \quad (3.17)$$

However,

$$\frac{\partial^2 \tilde{S}_{eff}}{\partial u_1^2} (u_1) = 0 \quad (3.18)$$

and

$$\frac{\partial^3 \tilde{S}_{eff}}{\partial u_1^3} (u_1) = -|b| \quad u_1 < 0 \quad (3.19)$$

so that (3.17) is not a stable minimum. The system evolves to $|u_1| = 1/2$ at which value it makes a transition to phase A_1 , *if* it has not already made this transition at a lower temperature. [See the discussion for (ii) above.]

(iv) $m_1^2 \leq 0$; $D < 0$

The solution to (3.5) is a single maximum at $u_1 < 0$, and not a minimum. There is no stable saddle-point for $|u_1| \leq 1/2$, so that at $T = T_H$ (*i.e.*, $m_1^2 = 0$), the system is already in phase A_1 .

In summary, for $(N_f/N) \sim \mathcal{O}(\lambda^2)$ and T sufficiently small, the system is in phase A_0 . At some $T_c < T_H$ there is a third-order phase transition to the high temperature phase A_1 . The free energy in phase A_0 behaves as

$$F/T \sim N_f^2 f_1(T) \quad 0 \leq T \leq T_c \quad (3.20)$$

and in phase A_1

$$F/T \sim N^2 f_2 \left(\frac{N_f}{N}, T \right) \quad T \geq T_c . \quad (3.21)$$

The low-temperature phase is interpreted as a gas of (color singlet) mesons and glueballs, which makes a third-order confinement-deconfinement transition to a gas of gluons and fundamental and anti-fundamental matter states at high temperatures.

4 Generalization

We use the strategy of ref. [8] to discuss the issues of this paper suitable for all orders of perturbation theory. In doing so, we modify their set-up to suit our problem. Recall that $u_n = 1/N \text{Tr } U^n$, so that (2.5) is a special case of

$$\begin{aligned}
S(U, U^+) &= \sum_{n=1}^{\infty} \frac{a_n(x)}{n} (Tr U^n)(Tr U^{+n}) \\
&+ N_f \sum_{n=1}^{\infty} \left[\frac{b_n(x)}{n} (Tr U^n) + c.c. \right] \\
&+ N^2 \sum_{\bar{k}, \bar{k}'} \alpha_{\bar{k}, \bar{k}'}(x) Y_{\bar{k}}(U) Y_{\bar{k}'}(U^+) \\
&+ N \left(\frac{N_f}{N} \right) \left[\sum_{\bar{k}, \bar{k}'} \beta_{\bar{k}, \bar{k}'}(x) Y_{\bar{k}}(U) Y_{\bar{k}'}(U^+) \right] \sum_{n=1}^{\infty} \left[(Tr U^n) + (Tr U^{+n}) \right] \quad (4.1)
\end{aligned}$$

where

$$Y_{\bar{k}}(U) = \prod_{j=0}^{\infty} (Tr(U/N)^j)^{k_j} \quad (4.2)$$

and where $\bar{k}$ and $\bar{k}'$ are arbitrary vectors of non-negative entries. Reality of the action and perturbation theory require [8]

$$\alpha_{\bar{k}, \bar{k}'} = \alpha_{\bar{k}', \bar{k}}^* = \alpha_{\bar{k}', \bar{k}} \quad (4.3)$$

and similarly for $\beta_{\bar{k}, \bar{k}'}$. It is useful to define

$$|\bar{k}| = \sum_j k_j \quad (4.4)$$

There is considerable redundancy in (4.1) which we remove in part by setting

$$\begin{aligned}
\alpha_{\bar{k}, \bar{k}} &= 0 \quad \text{for } |\bar{k}| = 1 \\
\alpha_{\bar{k}, 0} &= 0 \quad \text{for } |\bar{k}| = 1 \\
\beta_{0, 0} &= 0
\end{aligned} \quad (4.5)$$

Contact can be made with the formulation of (2.5) by noting that the leading behavior in perturbation theory of the coefficients in (3.1) are

$$\begin{aligned}
a_n(x) &= \mathcal{O}(1) \\
b_n(x) &= \mathcal{O}(1) \\
\alpha_{\bar{k}, \bar{k}'} &\sim (\lambda)^{|\bar{k}| + |\bar{k}'| - 2} \\
\beta_{\bar{k}, \bar{k}'} &\sim (\lambda)^{|\bar{k}| + |\bar{k}'|} .
\end{aligned} \quad (4.6)$$

Beyond the leading order of (4.6), the coefficients are a power series in λ and (N_f/N) , as well as any other coupling constants present in the problem.

We take advantage of the Gaussian trick extensively exploited in ref. [8] to write

$$\begin{aligned}
& \exp \left\{ \sum_{n=1}^p \frac{a_n}{n} \text{tr}(U^n) \text{tr}(U^{+n}) \right. \\
& \quad \left. + N_f \sum_{n=1}^p \frac{b_n}{n} [\text{tr}(U^n) + \text{tr}(U^{+n})] \right\} \\
& = \left(\frac{N^2}{2\pi} \right)^p \int \prod_{n=1}^p \left(\frac{a_n}{n} \right) dg_n d\bar{g}_n \exp -N^2 \sum_{n=1}^p \left(\frac{a_n}{n} \right) g_n \bar{g}_n \\
& \quad \cdot \exp N \sum_{n=1}^p \frac{1}{2n} [c_n \text{tr}(U^n) + \bar{c}_n \text{tr}(U^{+n})]
\end{aligned} \tag{4.7}$$

where

$$c_n = 2 \left[a_n g_n + \frac{N_f}{N} b_n \right]. \tag{4.8}$$

Eq'ns. (4.7)–(4.8) are closely related to (3.1) and (3.5) of ref. [5], except that now we allow a_n and b_n to be power series in (N_f/N) and λ , with the leading order of a_n and b_n of $\mathcal{O}(1)$, given by the zero-coupling expressions of (2.1).

Define the free-energy $F(g_n, \bar{g}_n)$ by

$$\begin{aligned}
& \exp [N^2 F(g_n, \bar{g}_n)] \\
& = \int [dU] \exp \left\{ N \sum_{n=1}^{\infty} \frac{c_n}{2n} [\text{tr}(U^n) + \text{tr}(U^{+n})] \right\}
\end{aligned} \tag{4.9}$$

which implies that

$$F(g_n, \bar{g}_n) = \frac{1}{4} \sum_{n=1}^{\infty} \frac{1}{n} c_n \bar{c}_n \tag{4.10}$$

in phase A_0 .

A. The Free-theory Revisited

It is instructive to consider further (4.1), but first with $\alpha_{\bar{k}, \bar{k}'} = \beta_{\bar{k}, \bar{k}'} = 0$. Using the Gaussian identity [8]

$$\begin{aligned}
& \exp \left[-N^2 \sum_{n=1}^p \left(\frac{a_n}{n} \right) g_n \bar{g}_n \right] \\
& = \left(\frac{N^2}{\pi} \right)^p \int \prod_{n=1}^p \left(\frac{a_n}{n} \right) d\mu_n d\bar{\mu}_n \\
& \quad \cdot \exp \left\{ -N^2 \sum_{n=1}^p \left(\frac{a_n}{n} \right) \mu_n \bar{\mu}_n + iN^2 \left(\frac{a_n}{n} \right) [\mu_n \bar{g}_n + \bar{\mu}_n g_n] \right\}.
\end{aligned} \tag{4.11}$$

The partition function is

$$\mathbb{Z} = \left(\frac{N^4}{2\pi^2} \right)^p \int \prod_{n=1}^p dg_n d\bar{g}_n d\mu_n d\bar{\mu}_n \exp(N^2 S_{eff}) \tag{4.12}$$

where here

$$S_{eff} = - \sum_{n=1}^p \left(\frac{a_n}{n} \right) \mu_n \bar{\mu}_n + i \sum_{n=1}^p \left(\frac{a_n}{n} \right) [\mu_n \bar{g}_n + \bar{\mu}_n g_n] + F(g_k, \bar{g}_k) . \quad (4.13)$$

The saddle-point equations are

$$\frac{\partial S_{eff}}{\partial g_k} = i \bar{\mu}_k + \left[a_k \bar{g}_k + \left(\frac{N_f}{N} \right) b_k \right] = 0 ; \quad (4.14)$$

$$\frac{\partial S_{eff}}{\partial \mu_k} = - \left(\frac{a_k}{k} \right) [\bar{\mu}_k + i \bar{g}_k] = 0 \quad (4.15)$$

and two analogous equations for variations with respect to $\bar{g}_k$ and $\bar{\mu}_k$. Since $a_k(x) \neq 0$, one has the solution

$$g_k = \bar{g}_k = \frac{(N_f/N) b_k}{(1 - a_k)} \quad (4.16)$$

since $a_k(x)$ and $b_k(x)$ are real. The density of eigenvalues in the phase A_0 is

$$\rho(\theta) = \frac{1}{2\pi} \left\{ 1 + \sum_{k=1} \left[g_k \exp(ik\theta) + \bar{g}_k \exp(-i k\theta) \right] \right\} , \quad (4.17)$$

so that from (4.8) and (4.16), one has

$$\rho_k = g_k = c_k/2 . \quad (4.18)$$

Thus, (4.16) is identical to eq'ns. (3.9) of ref. [5], where there is a 3rd order GWW phase-transition [13] to phase A_1 , interpreted as a confinement-deconfinement transition. Note that since $(N_f/N) \neq 0$, $g_k = 0$ is *not* a possible solution, so that the phase $\rho(\theta) = \text{constant}$ does not occur. Further discussion of the consequences of (4.9) and computation of $\rho(\theta)$ for phase A_1 may be found in ref. [5].

Thus, we have verified that the above formalism, with $\alpha_{\bar{k}, \bar{k}'} = \beta_{\bar{k}, \bar{k}'} = 0$, reproduces ref. [5]. We now return to the general case described by eq'ns. (4.1)–(4.6).

A. The General Case

Equation (4.13) generalizes to

$$\begin{aligned} S_{eff} = & - \sum_{n=1} \left(\frac{a_n}{n} \right) \mu_n \bar{\mu}_n \\ & + i \sum_{n=1} \left(\frac{a_n}{n} \right) [\mu_n \bar{g}_n + \bar{\mu}_n g_n] - F(g_k, \bar{g}_k) \\ & + \sum_{\bar{k}, \bar{k}'} \alpha_{\bar{k}, \bar{k}'} (-i)^{|\bar{k}| + |\bar{k}'|} Y_{\bar{k}}(\bar{\mu}) Y_{\bar{k}'}(\mu) \\ & + \left(\frac{N_f}{N} \right) \sum_{\bar{k}, \bar{k}'} \beta_{\bar{k}, \bar{k}'} (-i)^{|\bar{k}| + |\bar{k}'|} Y_{\bar{k}}(\bar{\mu}) Y_{\bar{k}'}(\mu) \sum_{n=1} (-i)^n [\mu_n + \bar{\mu}_n] \end{aligned} \quad (4.19)$$

where

$$Y_{\bar{k}}(\bar{\mu}) = \sum_{j=1} (\bar{\mu}_j)^{k_j}$$

and

$$Y_{\bar{k}'}(\mu) = \sum_{j=1} (\mu_j)^{k'_j} \quad (4.20)$$

The saddle-point equations at large N that follow from (4.19) are

$$\frac{\partial S_{eff}}{\partial g_k} = \left(\frac{a_k}{k} \right) \left\{ i \bar{\mu}_k + \left[a_k \bar{g}_k + \left(\frac{N_f}{N} \right) b_k \right] \right\} = 0 \quad (4.21)$$

$$\begin{aligned} \frac{\partial S_{eff}}{\partial \mu_j} &= - \left(\frac{a_j}{j} \right) [\bar{\mu}_j + i \bar{g}_j] \\ &+ \sum_{\bar{k}, \bar{k}'} \alpha_{k, k'} (-i)^{|k|+|k'|} \frac{(k'_j)}{\mu_j} Y_{\bar{k}}(\bar{\mu}) Y_{\bar{k}'}(\mu) \\ &+ \left(\frac{N_f}{N} \right) \sum_{\bar{k}, \bar{k}'} \beta_{\bar{k}, \bar{k}'} (-i)^{|\bar{k}|+|\bar{k}'|} \frac{(k'_j)}{\mu_j} Y_{\bar{k}}(\bar{\mu}) Y_{\bar{k}'}(\mu) \sum_{n=1} (-i)^n (\mu_n + \bar{\mu}_n) \\ &+ \left(\frac{N_f}{N} \right) \sum_{\bar{k}, \bar{k}'} \beta_{k, k'} (-i)^{|\bar{k}|+|\bar{k}'|} Y_{\bar{k}}(\bar{\mu}) Y_{\bar{k}'}(\mu) (-i)^j = 0 \end{aligned} \quad (4.22)$$

and two analogous equations for $\partial S_{eff}/\partial \bar{g}_j$ and $\partial S_{eff}/\partial \bar{\mu}_j$. Since $a_j \neq 0$, (4.21) implies that

$$\bar{\mu}_j = i \left[a_j \bar{g}_j + \left(\frac{N_f}{N} \right) b_j \right], \quad (4.23)$$

and similarly $\mu_j = i[a_j g_j + (N_f/N)b_j]$. One then substitutes (4.23) into (4.22) to obtain an equation for $\bar{g}_j$ in terms of a_j , b_j , $\alpha_{\bar{k}, \bar{k}'}$ and $\beta_{\bar{k}, \bar{k}'}$, and analogously for g_j . The solution to (4.22)–(4.23) for $\bar{g}_j$ and its analogue for g_j determines the distribution $\rho(\theta)$ in the ungapped phase, via (4.17) which is still valid.

We now ask whether $\rho(\theta) = \text{constant}$ is possible when $(N_f/N) \neq 0$. If so, this would require that $\bar{g}_j = 0$ for all j in (4.22)–(4.23), and analogously for g_j . If $k'_j \neq 0$, $g_j = \bar{g}_j = 0$ would require

$$\begin{aligned} \left(\frac{a_j}{j} \right) b_j &\stackrel{?}{=} \left\{ \sum_{|\bar{k}|, |\bar{k}'|} \alpha_{\bar{k}, \bar{k}'} \left(\frac{k'_j}{b_j} \right) \left(\frac{N_f}{N} \right)^{|\bar{k}|+|\bar{k}'|-2} \right. \\ &- 2 \sum_{|\bar{k}|, |\bar{k}'|} \beta_{\bar{k}, \bar{k}'} \left(\frac{k'_j}{b_j} \right) \left(\frac{N_f}{N} \right)^{|\bar{k}|+|\bar{k}'|} \left[\sum_n (-1)^{n+1} b_n \right] \\ &\left. - i \sum_{|\bar{k}|, |\bar{k}'|} \beta_{\bar{k}, \bar{k}'} \left(\frac{N_f}{N} \right)^{|\bar{k}|+|\bar{k}'|} (-i)^{j+1} \right\} Y_{\bar{k}}(b) Y_{\bar{k}'}(b). \end{aligned} \quad (4.24)$$

Note that a_j and b_j are both non-zero and of $\mathcal{O}(1)$ in (N_f/N) , while the right-hand side of (4.24) is a power series in (N_f/N) , beginning with leading terms of $\mathcal{O}(N_f/N)$ [see (4.5)]. Thus to any finite order in (N_f/N) (4.24) can not be satisfied by matching powers of (N_f/N) . One concludes that

$\rho(\theta) = \text{constant}$ is excluded in any finite order in the double perturbation theory. At low temperatures the system is in phase A_0 , and is *not* described by $\rho(\theta) = \text{constant}$ at any temperature.

The transition from phase A_0 to A_1 is driven by the third-order phase transition in $F(g_k, \bar{g}_k)$, *c.f.* (4.10) and (4.13), which occurs when $\rho(\pi) = 0$. This occurs when

$$g_1 = \bar{g}_1 \simeq 1/2. \quad (4.25)$$

[Equality occurs if one may neglect g_k for $k \geq 2$. See Sec. 3 for example.] Note that $a_j(x)$ and $b_j(x)$ are continuous functions of x as one passes through the (A_0, A_1) phase transition. From (4.23) this implies that $\bar{\mu}_j$ and $\bar{g}_j(x)$ [and similarly $\mu_j(x)$ and $g_j(x)$] are smoothly related to each other at the (A_0, A_1) boundary. Equation (4.19) can be written as

$$S_{eff} = -F(g_k, \bar{g}_k) + f(g_k, \bar{g}_k) \quad (4.26)$$

where $F(g_k, \bar{g}_k)$ has a third-order transition at the (A_0, A_1) boundary. Therefore it is plausible to assume that $f(g_k, \bar{g}_k)$ is smooth, with continuous derivatives at the phase-boundary. In the absence of an “accidental” third-order transition in $f(g_k, \bar{g}_k)$ which cancels that of $F(g_k, \bar{g}_k)$, or a lower-order discontinuity which seems unlikely given the smooth structure of $f(g_k, \bar{g}_k)$, S_{eff} has a third-order phase transition at the (A_0, A_1) boundary.

5 Conclusions

The phases of large N gauge theory at finite temperature on $S^{d-1} \times \text{time}$ are described by $\rho(\theta)$, the density of the eigenvalues of U in the unitary matrix model. When (N_f/N) is finite in the large N limit, $\rho(\theta) = \text{constant}$ is *not* a solution of the saddle-point equations at any finite order in the double perturbation theory. The saddle-point equations describe a phase A_0 , where $\rho(\theta) \neq \text{constant}$, but covers the interval $0 \leq \theta < 2\pi$. There is a transition to a second phase A_1 , where $\rho(\theta)$ is single gapped; the transition taking place at a critical temperature T_c , which is lower than a putative Hagedorn temperature. The above conclusions are correct to any finite order in a double perturbation series in the 't Hooft coupling λ and in (N_f/N) . In order for the low-temperature phase *not* to be A_0 , one presumably would require a phase-transition as one increased (N_f/N) at fixed λ , or increased λ at fixed (N_f/N) , keeping the temperature fixed. We speculate that this is not the case.

It is a reasonable assumption that generically $f(g_k, \bar{g}_k)$, *c.f.* (4.26), is sufficiently smooth that the (A_0, A_1) phase transition is controlled by $F(g_k, \bar{g}_k)$, *c.f.* (4.10) and (4.26), and is therefore 3rd order. It is plausible that these results, valid in any finite order of the double perturbation theory, should also hold at a strong coupling λ satisfying $N_f/N < 1/\lambda$ as well. Gravitational approximations are valid at large $g_s N$, so that the backreaction on D branes is small. To keep open string perturbation theory in control, one also needs $g_s N_f \ll 1$ which means $N_f/N \ll 1/\lambda$ in the large N limit. We conjecture that the third order phase transition is valid if these conditions are met.² This would then imply that the string theory dual to the gauge theory would exhibit a 3rd order Hawking–Page transition (dual to the confinement/deconfinement transition) in the supergravity limit of the string theory. No such transition has yet been found when $(N_f/N) = \text{finite}$ in the large N limit, as one must go beyond the probe approximation, and consider the

²We thank the referee for this argument.

back-reaction of the flavor branes. There have been preliminary studies involving the back-reaction of flavor branes [10, 11], but are not able to discuss the Hawking–Page phase transition from a thermal ensemble to a large black hole. Discovery of such a 3rd order transition in the string dual to a large N gauge theory with $(N_f/N) \neq 0$, would be an exciting confirmation of duality in the context of the ideas of this paper.

Acknowledgement

We thank Albion Lawrence for his comments and for reading the manuscript, and the referee for important suggestions for improving the paper.

References

- [1] E. Witten, Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150]. E. Witten, “Anti-de Sitter space, thermal phase transition, and confinements in gauge theories,” Adv. Theor. Math. Phys. **2**, 505 (1998) [arXiv:hep-th/9803131].
- [2] B. Sundborg, “The Hagedorn transition, deconfinement and N=4 SYM theory,” Nucl. Phys. **B573**, 349 (2000) [arXiv:hep-th/9908001].
- [3] O. Aharony, J. Marsano, S. Minwalla, K. Papadodimas and M. Van Raamsdonk, “The Hagedorn/deconfinement phase transition in weakly coupled large N gauge theories,” Adv. Theor. Math. Phys. **8**, 603 (2004) [arXiv:hep-th/0310285]; S. Minwalla and T. Wiseman, “Black hole-black string phase transitions in thermal 1+1 dimensional supersymmetric Yang–Mills theory on a circle,” Class. Quant. Grav. **21**, 5169 [arXiv:hep-th/0406210]; O. Aharony, J. Marsano, S. Minwalla, K. Papadodimas, M. Van Raamsdonk and T. Wiseman, “The phase structure of low dimensional large N gauge theories on tori,” JHEP **0601**, 140 (2006) [arXiv:hep-th/0508077].
- [4] O. Aharony, J. Marsano, S. Minwalla, K. Papadodimas and M. Van Raamsdonk, “A first order deconfinement transition in large N Yang–Mills theory on a small S^3 ,” Phys. Rev. **D71**, 125018 (2005) [arXiv:hep-th/0502149].
- [5] H.J. Schnitzer, “Confinement/deconfinement transition of large N gauge theories with N_f fundamentals: N_f/N finite,” Nucl. Phys. B695, 267 (2004) [arXiv:hep-th/0402219].
- [6] G.W. Semenoff, O. Tirkkonen and K. Zarembo, “Exact solution of the one-dimensional non-Abelian Coulomb gas at large N,” Phys. Rev. Lett. **D77**, 2174 (1996) [arXiv:hep-th/9605172]; G.W. Semenoff and K. Zarembo, “Adjoint non-Abelian Coulomb gas at large N,” Nucl. Phys. **B480**, 317 (1996) [arXiv:hep-th/9606117]; G.W. Semenoff, “Matrix model thermodynamics,” [arXiv:hep-th/0405107]; A. Dumitru, J. Leneghan, and D. Pisarski, “Deconfinement in matrix models about the Gross–Witten point,” [arXiv:hep-th/0410294]; P. Basu and S.R. Wadia, “R-charged AdS(5) black holes and large N unitary matrix models,” Phys. Rev. **D3**, 045022 (2006) [arXiv:hep-th/0506203]; D. Yamada and L.G. Yaffe, “Phase diagram of N=4 super-Yang–Mills theory with R-symmetry chemical potentials,” JHEP **0609**, 027 (2006) [arXiv:hep-th/0602074]; A. Karch and A. O’Bannon, “Chiral transition of N=4 Super Yang–Mills with flavor on a 3-sphere,” [arXiv:hep-th/0605120]; T. Harmark and M. Orselli, “Quantum mechanical sectors in thermal N=4 super Yang–Mills on $R \times S^3$,” [arXiv:hep-th/0605234]; “Matching the

- Hagedorn temperature in AdS/CFT,” [arXiv:hep-th/0608115]; C. Herzog, “A holographic prediction of the deconfinement temperature,” [arXiv:hep-th/0608151]; Y. Hikida, “Phase transitions of large N orbifold gauge theories,” [arXiv:hep-th/0610119]; T. Harmark, K.R. Kristjansson and M. Orselli, “Magnetic Heisenberg-chain/pp-wave correspondence,” [arXiv:hep-th/0611242]; “The Hagedorn temperature in a decoupled sector of AdS/CFT,” [arXiv:hep-th/0701088]; M. Brigante, G. Festuccia and H. Liu, “Hagedorn divergences in tachyon potential,” [arXiv:hep-th/0701205].
- [7] H. Liu, “Fine structure of Hagedorn transitions,” [arXiv:hep-th/0408001]; L. Alvarez-Gaumé, C. Gomez, H. Liu and S. Wadia, “Finite temperature effective action, AdS(5) black holes, and $1/N$ expansion,” Phys. Rev. **D71**, 124023 (2005) [arXiv:hep-th/0502227].
- [8] L. Alvarez-Gaumé, P. Basu, M. Mariño and S. Wadia, “Blackhole/string transition for the small Schwarzschild blackhole of AdS(5) $\times$ S⁵ and critical unitary matrix models,” [arXiv:hep-th/0605041].
- [9] S.W. Hawking and D.N. Page, “Thermodynamics of black holes in anti-de Sitter space,” Comm. Math. Phys. **87**, 577 (1983); A. Buchel and L.A. Pando Zayas, “Hagedorn vs. Hawking-Page transition in string theory,” [arXiv:hep-th/0305179].
- [10] S. Cherkis and A. Hashimoto, “Supergravity solution of intersecting branes and AdS/CFT with flavor,” JHEP **0211**, 036 (2002) [arXiv:hep-th/0210105]; M. Gómez-Reino, S. Naculich and H. Schnitzer, “Thermodynamics of the localized D2-D6 system,” Nucl. Phys. **B713**, 263 (2005) [arXiv:hep-th/0412015].
- [11] R. Casero, C. Núñez and A. Paredes, “Toward the string dual of $N=1$ SQCD-like theories,” Phys. Rev. **D73**, 086005 (2006) [arXiv:hep-th/0602027]; A. Paredes, “On unquenched $N=2$ holographic flavor,” [arXiv:hep-th/0610270]; B. Burrington, J.T. Liu, L. Pando-Zayas and D. Vaman, “Holographic duals of flavored $\mathcal{N}=1$ super Yang–Mills: Beyond the probe approximation,” [arXiv:hep-th/0406207]; J. Babington, J. Erdmenger, N. Evans, Z. Guralnik and I. Kirsch, “Chiral symmetry breaking and pions in non-supersymmetric gauge/gravity duals,” Phys. Rev. **D69**, 066007 (2004), [arXiv:hep-th/0306018]; I. Kirsch, “Generalizations of the AdS/CFT correspondence,” [arXiv:hep-th/0406274]; J. Erdmenger and I. Kirsch, “Mesons in gauge/gravity dual with large number of fundamental fields,” JHEP **0412**, 025 (2004) [arXiv:hep-th/0408113]; R. Apreda, J. Erdmenger, N. Evans and Z. Guralnik, “Strong coupling effective Higgs potential and a first order thermal phase transition from AdS/CFT duality,” Phys. Rev. **D71**, 126002 (2005) [arXiv:hep-th/0504151]; F. Benini, F. Canoura, S. Cremonesi, C. Núñez and A.V. Ramallo, “Unquenched flavors in the Klebanov–Witten model,” [arXiv:hep-th/0612118].
- [12] D. Mateos, R. Myers and R. Thomson, “Holographic phase transitions with fundamental matter,” [arXiv:hep-th/0605046]; “Thermodynamics of branes,” [arXiv:hep-th/0701132].
- [13] D. Gross and E. Witten, “Possible third order phase transition in the large N lattice gauge theory,” Phys. Rev. **D21**, 446 (1980); S. Wadia, “ $N=\infty$ phase transition in a class of exactly soluble model lattice gauge theories,” Phys. Lett. **B93**, 403 (1980); “A study of $U(N)$ lattice gauge theory in two-dimensions,” EFI-79/44-CHICAGO.
- [14] A. Karch and E. Katz, “Adding flavor to AdS/CFT,” JHEP **0206**, 043 (2002) [arXiv:hep-th/0205236].

- [15] J. Jurkiewicz and K. Zalewski, “Phase structure of $U(N \rightarrow \infty)$ gauge theory on a two-dimensional lattice for a broad class of variant actions,” Nucl. Phys. **B220** [FS8] 167 (1983).
- [16] M. Kruczinski and A. Lawrence, “Random walks and the Hagedorn transition,” JHEP **0607**, 031 (2006) [arXiv:hep-th/0508148].